

# Evaluation of Noise Pollution in Small Workshops in Qom, Iran; and Its Situation Compared to National Noise Standards

Akbar Scandari<sup>a</sup>, Hossein Jafari Mansoorian<sup>b</sup>, Alireza Mashkori<sup>c</sup>, Zahra Ahmadli<sup>a</sup>, Narges Khanjani<sup>d</sup>, Samira Norzaee<sup>e</sup>, Mohammad Fahiminia<sup>f\*</sup>

<sup>a</sup>Department of Environmental Health Engineering, Qom University of Medical of Sciences, Qom, Iran.

<sup>b</sup>Young Researchers and Elite Club, Hamedan Branch, Islamic Azad University, Hamedan, Iran.

<sup>c</sup>Department of Occupational Health, Faculty of Health, Qom University of Medical Sciences, Qom, Iran.

<sup>d</sup>Environmental Health Engineering Research Center, Kerman University of Medical Sciences, Kerman, Iran.

<sup>e</sup>Department of Environmental Health Engineering, Iranshahr University of Medical Sciences, Iranshahr, Iran.

<sup>f</sup>Research Center for Environmental Pollutants, Qom University of Medical Sciences, Qom, Iran.

\*Correspondence should be addressed to Dr. Mohammad Fahiminia, Email: [fahiminia.m@gmail.com](mailto:fahiminia.m@gmail.com)

## A-R-T-I-C-L-E-I-N-F-O

### Article Notes:

Received: Mar 8, 2018

Received in revised form:

Agu 20, 2018

Accepted: Sep 19, 2018

Available Online: Oct 7, 2018

### Keywords:

Noise pollution, noise, small workshops, industrial noise, Qom, Iran

## A-B-S-T-R-A-C-T

**Background & Aims of the Study:** Noise is one of the most important environmental pollutants. It is a global health problem, especially among industrial workers. Since there was little information about noise levels in small workshops in Qom city, this study aimed to noise evaluation in small workshops and compares it with national standards.

**Materials and Methods:** This was a cross-sectional study. Initially jobs and workshops that were causing noise pollution were identified. Then, noise was measured with a calibrated sound dosimeter, model CASELLA CEL-320, in A-network, in 108 working places. The results were compared with national noise standards.

**Results:** The total numbers of workers in the workshops were 317 people, and inside the workshops 76.3% of the workers and 66.7% of the workshops were exposed to noise over the standard threshold. Outside the workshops, in 89% noise pollution was higher than the standard city values. Only 11% of noise pollution in the city workshops had a 1 or less than 1 dB difference with the standard noise threshold.

**Conclusion:** Small workshops cause a significant amount of noise pollution in Qom city. This can affect the workers' and the public's health. Noise control procedures such as controlling the source and direction of noise distribution, as well as setting up workshops in the right location, and far from residential areas are necessary.

**Please cite this article as:** Scandari A, Jafari Mansoorian H, Mashkori A, Ahmadli Z, Khanjani N, Norzaee S, Fahiminia M. Evaluation of Noise Pollution in Small Workshops in Qom, Iran; and Its Situation Compared to National Noise Standards. Arch Hyg Sci 2018;7(3): 157-164

## Background

Population growth along with industrial and technological development in megacities has increased problems for urban dwellers and one of these problems is environmental pollution (1). One of the main environmental problems that have been less studied in Iran is noise pollution (2), which is now a global health problem, especially for industrial workers (3).

Although the industrial revolution caused increased growth in production in the world, but it has also made noise an inevitable exposure for humans (4).

Noise pollution is an important and common health hazard in most industries, and if necessary preventive measures are not implemented it can lead to physiologic, mental, economic and social problems among the exposed employees. Noise above the stranded

threshold leads to hearing loss, nuisance in conversation, distraction and not noticing danger signs; and therefore adversely affects workers safety and performance. It is also causes stress and unwanted physiological responses and decreased comfort among workers during work time. Although evaluating and controlling noise in big industrial environments has been considered by researchers, but in small scale industries due to various reasons, noise evaluation has been taken less seriously (5,6).

Small workshops create jobs for urban populations, have a close link with other parts of the society and are important in urban development. However, the presence of these industries and workshops especially in residential areas can increase pollution and exposure to health hazards, such as noise, waste, particulate matter, bad orders, vibration, wastewater, soot and air pollution. These industries might also cause traffic and nuisance (7,8).

It has been estimated that in developing countries, 45 to 95% of the workforce work in small scale industries (9). Small scale industries according to the definition of the Iran Statistics Centre are industries in which the number of employees is not more than 9 people. Statistics reported by the Iran Statistics Centre in 1987 shows that from 339234 industrial workshops inside the country a number of 327715 which is about 96.5 % of the active workshops, are small workshops with 1 to 50 employees and these workshops make 64.5% of industrial occupations (10). Also according to statistics reported by the Iran Industrial Organization from 89991 industrial licenses granted by the end of 2011, 80289 cases, which are 89.2%, were granted to workshops with less than 50 employees (11). The occupation rate in Qom province in 2006 was 94.2% which means the industrial and services sector in this province has a good situation and respectively 42.6 and 49.2% of all occupations were in this sector (12). Noise pollution from industrial activity in

Qom city, along with noise from traffic is among the problems of people residing close to industries, workshops and roads.

Noise pollution is one of the most prevalent occupational hazardous risk factors; and can have adverse effects on different organs such as the hearing and cardiovascular system. It also has negative effects on mental health and work efficiency among workers.

#### **Aims of the study:**

In this study we tried to estimate the level of noise pollution in small workshops of Qom city and compare it with occupational standards and then suggest ways for controlling noise pollution.

#### **Materials & Methods**

This was a cross-sectional descriptive study conducted in May, June and July of 2015 in order to measure noise pollution in the small workshops of Qom City which had less than 10 employees. Initially a list of all private occupations was inquired from the Guild Hall of Qom City. Occupations with workshops which were causing noise pollution were identified, and then 108 workshops, and from each workshop three cases were randomly selected. The total number of workers in the selected workshops was 317 people. Measuring noise was done for 30 minutes inside the workshops; and also 30 minutes in a one meter distance outside the workshop at the peak of their occupational activity in daytime and at two times, one in midday (10:30-13:30) and one in afternoon (16:30-19:30) hours. Because noise intensity is on a logarithmic scale, in order to estimate the average of noise levels in three workshops, equation number 1 was used. Also this formula was used to estimate the average of noise levels in three spots which were in the commuting areas of the workers and in workshops with pounding noises. This equation is used for multi-levels in different frequencies and different sources for one level, and calculates the overall sound level. Due to the logarithmic nature of sound pressure levels,

it is not possible to directly do summation, subtraction or averaging on these numbers. After voice measurements, in order to compare the results with the threshold, the average of time levels (SPLTWA) was calculated for an 8-hour shift, by using equation 2.

In calculating the average of time levels, first the level of each exposure and the corresponding exposure time is measured, then the equivalent exposure levels of the worker was calculated for a period of time according to the equation. After calculating the equivalent level of occupational exposure, the results were compared with the standard of the professional occupational health committee of Iran, which is 85 dB for a working shift (8 hours). As the area understudy was generally among the residential industrial areas, in this study we compared the sound pressure levels outside the workshops with the standard for these areas which is 70 decibels (13).

$$LP (dc) = 10\text{Log} \left[ \frac{1}{n} \sum_{i=1}^n 10 \frac{LP_i}{10} \right] \quad \text{Eq. (1)}$$

Where LP is the average source sound intensity, N is the number of measuring points and LP<sub>i</sub> is the sound pressure level in each measuring point.

$$Leq (db.) = 10\text{Log} \left[ \frac{1}{T} \sum_{i=1}^n ti 10 \frac{LP_i}{10} \right] \quad \text{Eq. (2)}$$

Where Leq is sound level equilibrium in decibels, ti is the duration of exposure in hours, T is the reference time of 8 hours and LP<sub>i</sub> is the sound pressure level of exposure in decibels.

### Noise measuring

Noise measuring was done by local measurements in order to estimate workers exposure and by using a sound dosimeter, model 320-CEL made in England by CASELLA with a CEL-282 caliber and through these steps.

1. The dosimeter was calibrated through the method advised by the company and the A-

network was used for measuring general sound levels. The device was calibrated after each measurement in a quiet environment.

2. Sound measurement was done according to the guidelines of the Occupational Health Technical Committee of Iran.

3. The speed of the device was chosen as slow according to standard recommendations.

4. The duration of measuring sound was 30 minutes inside the workshop and 30 minutes in a 1 meter distance outside the workshop.

5. The variables measured by the device including: LAS, LAS<sub>mn</sub>, LCPK, LAE and LCPK were recorded.

### Results

Table 1 shows the average number of employees in each workshop, the average measured sound pressure levels (LAEQ) inside the workshops and 1 meter outside the workshops, the time average of the levels (SPLTWA) and the average working hours of the workers in a day. The levels measured inside the workshops are the sound level which workers are exposed to during working hours. The main source of noise production in these workshops was the sharpening machine, hammer, trunk mold, wind pump, injector engine speed testing machine, firesaw, cupboard cutting machines, three-functioning saws, milling machines, motorcycle testing and tuning, knitting machines and hosiery making machines, sewing machines, whipping machines (in carpet washing), offset machine (in printing shop), printing services machines, milling and peaking, grinding, and cutting machines and confectionary mixer machines. The noise recorded 1 meter away of the workshops, was from the activities of neighboring workshops and commuting vehicles, as well as the noise from the workshop machines.

**Table 1) Noise from machinery and workshop activities (LAEQ) in different occupations**

Occupational group	activity	Number of units	Average number of employees	Average working hours per day	Leq (30min) inside the workshop	SPL <sub>TWA</sub>	LEQ (30 min) in a 1 m distance outside the workshop
<b>Aluminium makers</b>	Aluminium door and window frame	3	2.33	8	98.4	98.4	79.4
<b>Car dealers</b>	Car dealer representative branches and mechanics	3	11	9	96.2	96.7	85.7
<b>Shirts and trico</b>	Shirt maker	3	1	7	74.4	73.8	69
<b>Blacksmiths and cutters</b>	Lathing	3	2.66	10	92.2	93.1	81.1
	Car Blacksmithing	3	2	8	87.9	87.9	78.5
	Cylinder making	3	2.66	7	89.4	88.8	85
	Welding water pump pieces	3	1.33	7	77.3	76.7	73.2
<b>Car oil change</b>	Changing car oil and air	3	3	10	94.1	95	81.1
	Carwashing	3	5	10	86.6	87.7	76.1
<b>Rag making</b>	Rug washing	3	6	12	88.2	89.9	69.7
<b>Printing and binding</b>	Printing	3	4	11	81.5	82.8	73.3
	Binding	3	2	10	87.9	88.8	73.8
	Printing services	3	2	10	76.5	77.4	69.8
<b>Tailors</b>	Tailors	3	1.66	9	74.4	74.9	69
<b>Blacksmiths and window frame makers</b>	Welding	3	1.33	8	77.3	77.3	73.2
	Iron doors and windows	3	3	10	102.8	103.8	93.3
<b>Carpenter</b>	Wood cutting	3	3.33	9	102.5	103	75.1
	Wardrobe building	3	2	11	84	85.4	74.1
	Furniture making	3	2.33	11	82.4	83.8	76.3
	Carpentry	3	2	8	93.1	93.1	80.5
<b>Bicycle and motor cycle</b>	Motor cycle repairing	3	1	4	89.1	86	81
<b>Tile making</b>	Stone cutting	3	2.33	9	82.8	83.3	68.9
<b>Stone cutters</b>	Stone cutting services	3	3	11	93.4	94.8	86.7
<b>Flattening</b>	Car ironing	3	2	7	101	100.4	76.2
<b>Carpets and decorations</b>	Making blinds and curtains	3	2.33	11	90.9	92.3	74.8
<b>Metal-workers</b>	Canal making,	3	2	7	93.1	92.5	92.9
	Making cooler valves	3	3	19	85.9	86.8	76.1
<b>Locks, hinges and hardware</b>	Key making	3	1	6	81.6	80.3	74.3
<b>Confectioners</b>	Honey and Sesame Sohan	3	4.33	11	80.7	82.1	75.2
<b>Metal product producers</b>	Kitchen cupboard makers	3	2	10	96.2	97.1	88.5
<b>Knitter</b>	Knitting with computer devices,	3	3	9	87.8	88.3	74.5
	Sock making with computer devices	3	3	8	76.4	76.4	71.6
<b>Construction material and equipment</b>	Iron selling and Blacksmithing	3	2.33	10	88.4	89.4	80
<b>Mechanics</b>	Injector pump	3	2	8	88.3	88.3	69.7
	licensed mechanic	3	6.66	9	93.6	94.1	76.4
<b>Tile makers</b>	Tile making	3	3	10	80.6	81.6	76
<b>Total</b>	-	108	317	-	-	-	-




Figure.1) Comparing the noise inside the Workshops with National standards




Figure 2) Comparing sound levels outside the workshops with national standard thresholds

According to the results in Fig.1 and comparing it with national standards, all occupations related to the car industry produced more than

85 decibels noise and according to the time average levels in table 1 and the recommendations of the national occupational

health technical committee, in the aluminum door and window frame industry and the iron door and window frame industry, the situation is similar to workers in the automobile industry and workers in these groups are probably more prone to the adverse effects of high noises. Our results show that 76.34% of the workers in 66.67% of the workshops received noise levels over the standard threshold.

Results in Fig. 2 show that except a few workshops, other measurements done in a 1 meter distance from workshops were all above the national threshold for industrial residential areas. This threshold is 70 decibels and the workshops with lower than standard threshold were just a few decibels lower than the threshold. According to the results in fig. 2, in 89% of the measurements, noise pollution outside the workshop was above standard limits and in only 11%, it was in the standard range.

## Discussion

In this study, the average daily sound levels in most small workshops in Qom city was over the standard threshold. Also the sound level of more than 66.7% of the workshops, inside the workshop and 89% of the workshops in a 1 meter distance away from the workshop was higher than the permissible national standard.

In other studies conducted in Iran, sounds levels in the majority of small urban workshops was above standard thresholds as well. Rashidi *et al* (14) reported that in the towns of Ilam Province, blacksmiths were producing the highest level of noise pollution in their workshops with sound levels over 103.8 decibels; and then iron door and window frame makers with a sound level of 102.8 decibels has the second highest level of noise measured. In Rashidi's study noise pollution in the printing industry was 79.6 decibels. In our study the noise level was 81.54 dB in printing workshops, 87.9 dB in binding workshops and 76.5 decibels in printing services. In another study done by Hokm Ali *et al* (15) in small

workshops of Birjand City, the level of noise pollution in stone carving workshops was 94.5 decibels. In our study the level of noise pollution in stone carving workshops was 93.4 decibels. Also in a study done by Taheri *et al* noise levels in carpentry workshops were 92.8 dB, which is similar to the current study in which the levels in carpentry workshops were 93.1 dB, in furniture making 82.4 dB, in cupboard making 84 dB and in wood cutting 102.5 dB. The sound level at the car body repair workshop was equal to 100.4 dB. This value was similar to the sound measured in Qazvin city body repair workshops, which had an average workers daily exposure to noise of  $98.3 \pm 3.4$  dB in network A, and was in a range of 90-140 dB (17). Also, in Taheri *et al*'s study about the noise level of carpentry workshops in Sabzevar, noise exposure was 92.8 dB and the minimum and maximum were 72 and 115.3 dB respectively (18). Noise exposure in the present study was 103 for wood cutters, 85.4 for closet makers, 83.8 for furniture makers and 93.1 dB for carpenters. These results are in the range of noise pollution that the carpentry workshops of Sabzevar city had. In this current study the area was dominantly residential-industrial, and the sound levels outside the workshops was compared with the standards for these areas; and except in 11% of the workshops, the noise levels outside the workshops was above the standard levels for residential-industrial areas (70 dB).

High noise levels above the threshold have also been reported from other cities in Iran. In a study (19) about evaluating noise pollution in Qazvin city, noise levels in the city commuting areas was between 69.9 to 72.8 dB. In the study done by Naddafi *et al* (20) the maximum noise pollution in different parts of Zanjan during day and night was respectively 72 and 77.7 dB, in the A network and above the standard threshold. In the residential-industrial areas of Tabriz city, the amount of noise pollution in the studied areas was between 69.67 and 70.77 dB (21), which is higher than the standard noise

level for these areas. Other studies about noise pollution in cities has shown that in most city regions the noise levels are higher than national standards and factors such as industries and traffic are among the main reasons for noise pollution (22,23).

Other studies conducted outside Iran, such as a study done by Henrique et al (24) in the city of Curitiba in Brazil showed that in 93.3% of the measurements, the levels of noise pollution was more than 65 dB and in 40.3% noise pollution was more than 75 dB. Also in another study done by Yilmaz and Ozer (25) in 126 points in the city of Erzurum in Turkey, in 96 measuring points, noise levels were higher than standard. Darson et al (26) evaluated noise pollution in Ghoonieh, Turkey; and noise measurements were done in 366 points of the city and the average noise pollution of the city was above 65 dB; and the main sources of noise production were traffic and industries.

### Conclusion

Most small workshops in cities produce significant noise pollution which is higher than the recommended national standards. Due to the hazardous effects of noise on workers and people residing in these areas, we suggest that the basics of occupational safety should be taught to workers, protective equipment should be supplied to them, their working hours should decrease and their rest should increase, appropriate industrial equipment should be chosen for the production processes, equipment should be taken care of and serviced regularly, equipment should be left in the appropriate location and should have sound insulators or should been closed in confined insulating spaces. In cities, workshops should be established in proper locations and regulations should prevent founding workshops in residential areas. A program for organizing industries and businesses, based on the experience of other Iranian and world cities (27,

28) can be efficient in controlling noise pollution in Qom city.

### Footnotes

#### Conflict of Interest:

The authors declared no conflict of interest.

### References

1. Chopani M. Environmental pollutants and environmental protection: Publishers Manpower training and equipping of National Iranian Gas Company. 2009;(in Persian).
2. Bahram soltani K. Basics of Architectural urban green space: Department of Urban Planning and Architecture Department of Housing and Urban Development-Secretariat of the Supreme Council for Planning and Architecture of Iran2005. 64-108 (in Persian).
3. Karlidağ T, Yalçın Ş, Öztürk A, Üstündağ B, Gök Ü, Kaygusuz İ, et al. The role of free oxygen radicals in noise induced hearing loss: effects of melatonin and methylprednisolone. *Auris Nasus Larynx*. 2002;29(2):147-52 (in Persian).
4. Moharam njhad N, Rashidi A, Enayati VR. Evaluation of noise pollution in the spinning and weaving factory in Qazvin province and the city of Najafabad, and assess the negative effects on workers exposed. *Journal of Textile Technology*. 2011;6(1):22-8 (in Persian).
5. Aghilinejad M, Alimohamadi L, Mohammadi S, Fallahi M. Assessment of the effect of occupational noise on workers hearing in small scale industries in Tehran. *Journal of Military Medical University Islamic Republic of Iran*. 2007;5:1305-10 (in Persian).
6. Nasiri P, Farahng deghan S, Monazam M. Noise pollution assessment and feasibility control methods in a petrochemical plant. *Journal of Occupational Health and Safety*. 2011;1(1).
7. Garakhlo M, Fargam R. Accounting and optimal deployment of industry and urban disturb yards (Case Study: Central texture of Kermanshah). *Geographical research*. 2001;40:47-71 (in Persian).
8. Nasiri E. Small workshop industries and its impact on urban development productivity"The study of Bvhn". *Urban Regional Studies and Research*. 2009;1(1):129-42 (in Persian).
9. aghilinejad M, Mohammadi A, Mohammadi S, Falahi M. Assessment of the effect of occupational noise on workers hearing in small scale industries in Tehran. 2007;5(3):1305-10 (in Persian).
10. Miri A, Hagigi M. The effect of environmental factors on small business marketing plan Iran. *Knowledge management*. 1996;33:(in Persian).

11. Malakouti J, Rahimi H. Investigation heat stress in small enterprise in Qom city. *Journal of Health and Safety at Work*. 2014;3(4):59-68 (in Persian).
12. Iranjhad MH, Bashiri H, Bageri H, Naini MR, Srezai A, Fatahi MM, et al. Evaluation of the quality and quantity of green space the city of Qom and solutions to achieve optimal situation. The third national conference on green space and urban landscape. The National Conference of green space and urban landscape. 2008:88-98 (in Persian).
13. Golmohamadi R. Sound and vibration engineering in the industry and environment: Publishers daneshjo.himedan 2010 (in Persian).
14. Rashidi R. Evaluation of noise pollution and its influencing factors in the industries and occupations city of Ilam. Ilam University of Medical Science. 2001;89(29):36-44 (in Persian).
15. Hokm Abadi R, Amiri H, Nekooi N, Mozafarian S. Sixteenth National Congress of the Iran-Mehr Environmental Health. 2013:(in Persian).
16. Taheri m, Kalat Alabi H. Hearing loss assessment in Sabzevar carpenters. Procdings of the first Internashonal conferens of HSE 2008: Sep15-17 Isfahan, Iran (in Persian).
17. Ahmadi S, Karboro AA, Einanlo M, Aubi zade H, Zarei M. Occupational Noise Exposure and Hearing Loss Among Car Smoothers in Qazvin. *Health & Environ*. 2011;4(1):85-92 (in Persian).
18. Halvani Gh. Noise exposure and hearing loss assessment in taban yazd textile workers. *J. Birjand Medical University*.2008; 15(4):69-74(in Persian).
19. Emam jomee M, Nik pi A, Safari Varyani A. Study of noise pollution in Qazvin *Journal of Qazvin University of Medical Sciences*. 2011;15(1):64-70 (in Persian).
20. Naddafi K, Yunesian M, Mesdaghinia A, Mahvi A, Asgari A. Noise pollution in Zanjan city in 2007. *Zanjan University of Medical Sciences*. J. 2008;16(62):85-96(in Persian).
21. Ghanbari M, Naddafi K, Mosaferi M, Yunesian M, Aslani H. Noise Pollution Evaluation in Residential and Residential-Commercial Areas in Tabriz-Iran. *Iranian Journal of Health & Environment*. 2011;4(3):375-4 (in Persian).
22. Tajic R, Ghadami A, Ghamari F. The effects of Noise Pollution and Hearing of metal Workers in Arak. *Zahedan Journal of Research in Medical Sciences*. 2008;10(4):291-8 (in Persian).
23. Mari Oryad H, Raygan Shirazi A, Alimohammadi I. Evaluation of noise pollution in crowded areas in Yasuj in 2006. *Armaghane danesh*. 2008;12(4):109-16 (in Persian).
24. Zannin PHT, Diniz FB, Barbosa WA. Environmental noise pollution in the city of Curitiba, Brazil. *Applied Acoustics*. 2002;63(4):351-8.
25. Yilmaz H, Ozer S. Evaluation and analysis of environmental noise pollution in the city of Erzurum, Turkey. *International Journal of Environment and Pollution*. 2005;23(4):438-48.
26. Dursun S, Ozdemir C, Karabork H, Kocak S. Noise pollution and map of Konya city in Turkey. *Journal of International Environmental Application & Science*. 2006;1(1):63-72.
27. Rahimian S, Akhondzade HR, Nasr M, Katkhodazade Z. Business studies urban pollution and nuisance in Shiraz. Shiraz Municipality. URL:e-exhibition.tehran.ir/Article-View/ArticleId/2846 (in Persian).
28. Fatah RA, soltani F. Industry and unions organizing in cities: Publications of the country's municipalities Vdhyary; 2011 (in Persian).